

ORIGINAL ARTICLE

Epidemiology, Genetics & Prevention

The role of fish intake on asthma in children: A meta-analysis of observational studies

M. M. Papamichael¹  | S. K. Shrestha² | C. Itsiopoulos¹ | B. Erbas³ 

¹Department of Rehabilitation, Nutrition and Sport, School of Allied Health, La Trobe University, Melbourne, VIC, Australia

²Research Centre for Integrated Development (RECID), Nepal, Nepal

³School of Psychology and Public Health, La Trobe University, Melbourne, VIC, Australia

Correspondence

Maria Michelle Papamichael, La Trobe University, School of Allied Health, Dept of Rehabilitation, Nutrition & Sport, Melbourne, VIC, Australia.
Email: sassipap@hotmail.com

Abstract

Background: The evidence is mixed on the use of long chain Omega-3 fatty acids in the prevention and management of childhood asthma.

Methods: We conducted a systematic search and meta-analysis investigating the role of fish intake, the main dietary source of long chain omega-3 fatty acids, on asthma in children.

Results: A total of 1119 publications were identified. Twenty-three studies on fish intake in association with childhood asthma were included in the final review. In 15 of 23 studies, early introduction of fish (6-9 months) and regular consumption (at least once a week) improved asthma symptoms and reduced risk in children 0-14 years as compared to no fish consumption; 6 of 23 showed no effect and 2 of 23 studies suggest adverse effects. Meta-analysis revealed an overall "beneficial effect" for "all fish" intake on "current asthma" [OR: 0.75; 95%CI: 0.60-0.95] and "current wheeze" [OR: 0.62; 95%CI: 0.48-0.80] in children up to 4.5 years old. An overall protective effect of "fatty fish" intake as compared to "no fish" intake in children 8-14 years old was also observed [OR: 0.35; 95%CI: 0.18-0.67].

Conclusion: This meta-analysis suggests that introduction of fish early in life (6-9 months) and regular consumption of all fish (at least once a week) reduces asthma and wheeze in children up to 4.5 years old, while fatty fish intake may be beneficial in older children. Future well-designed clinical trials are recommended to confirm the promising findings documented in this literature analysis.

KEYWORDS

asthma, children, fish, nutrition, oily fish, Omega-3 fatty acids

1 | INTRODUCTION

Despite advances in pharmacotherapy, globally, the asthma epidemic in children is a major public health problem as it continues to rise. Asthma is associated with morbidity, substantial healthcare costs and decreased quality of life.^{1,2} There is no cure for asthma; symptoms can only be controlled by medication which on a long-term basis may cause side effects. Therefore, identifying potential non-pharmacological interventions which improve management of asthma is of great public health significance.

The modern diet is low in fruits, vegetables, fish and high in refined grains, red meat and fast food³ which may have contributed to the high prevalence of asthma in Westernized countries.¹ As a result today's diet is deficient in anti-inflammatory nutrients (from fruit/vegetables) and long-chain Omega-3 fatty acids (LC n-3 PUFA) (obtained from fish) and high in Omega-6 fatty acids mainly from processed fats and oils such as margarine and vegetable oils.^{3,4} It is believed that the change in anti-inflammatory status, fatty acid composition, increase in Omega-6: Omega-3 fatty acid ratio is associated with an increase in oxidative stress

and airway inflammation promoting the development and symptoms of asthma.^{5,6}

There has been considerable interest in the potential therapeutic and protective properties of Omega-3 fatty acids (EPA/DHA) in the pathogenesis of asthma due to anti-inflammatory and immunomodulating properties.^{4,7} To date, the evidence for the beneficial effects of Omega-3 fatty acid intake in asthma is controversial and intervention trials investigating the impact of fish or fish oil supplementation in children are lacking.⁸ Most of the epidemiological evidence which gave rise to the hypothesis that marine Omega-3 fatty acids may have a protective effect is based on studies documenting that regular consumption of fish has a prophylactic effect.⁹⁻¹¹ In addition, many systematic reviews have investigated the impact of Omega-3 fatty acid intake during pregnancy on allergy outcome in offspring suggesting a beneficial effect.¹²⁻¹⁴ Little is known about fish intake on asthma during childhood (post-infancy) and this warrants further investigation.

The purpose of this study is to conduct an up-to-date systematic search, qualitatively synthesize the available evidence and perform a meta-analysis to determine the role of fish (lean and fatty) on childhood asthma. The impact of fish intake during pregnancy is beyond the scope of this review.

2 | METHODS

This review was conducted according to PRISMA guidelines¹⁵ (Data S4).

2.1 | Literature search strategy

We conducted a comprehensive literature search of publications up to July 2017 using the following bibliographic databases: Cochrane Central Register of Controlled Trials, MEDLINE, PubMed, CINAHL (EBSCO), SCOPUS and EMBASE. No restrictions were applied on language or publication dates. Supplementary studies were sought from conference proceedings as well as clinical trials registries (International, European, Australia and New Zealand) in order to identify published/unpublished trials. Other potential relevant citations were identified by hand search of the reference lists of relevant articles, reviews, systematic reviews and meta-analysis. Databases were searched for relevant publications, by a two-step search strategy using the following search terms:

Search 1: ["fish OR fatty fish OR oily fish OR omega 3 fatty acids OR n-3 long chain polyunsaturated fatty acids AND "childhood asthma"]

Search 2: ["fish OR fatty fish OR oily fish OR omega 3 fatty acids OR n-3 long chain polyunsaturated fatty acids AND "children" AND "asthma"].

The final search was a combination of both searches using the term "OR." The full search strategy is provided in (Box S1).

2.2 | Study eligibility criteria

2.2.1 | Inclusion criteria

Inclusion criteria was based on Participants, Intervention, Comparator and Outcomes (PICO).¹⁶

2.2.2 | Type of participants

This systematic review considered publications that included children younger than 18 years old.

2.2.3 | Type of intervention

Publications reporting the effect of fish as the primary exposure measurement.

2.3 | Outcome measures

Primary outcomes of interest were all asthma symptoms (wheeze, dyspnoea, shortness of breath) and prevalence or incidence of asthma.

A publication was considered when exposures were measured and the presence of asthma was based on the manifestation of symptoms, pulmonary tests, or doctor-diagnosis.¹⁷ In addition, studies examining the effect of Omega-3 fatty acid intake on atopy in children were considered relevant when results on fish intake and asthma outcome in children could be separated.

2.4 | Type of studies

We considered experimental and epidemiological study designs including randomized controlled trials (RCTs), non-randomized controlled trials, before and after intervention studies, cohort, case-control and cross-sectional studies.

2.5 | Exclusion criteria

Exclusion criteria were based on PICO characteristics and reasons for exclusion are summarized in (Figure 1). Publications not included were as follows: reviews, systematic reviews, editorials, comments, letters, case studies, animal studies and those with no abstract, full text or English translation available. Interventions or risk factors other than fish consumption such as: formula supplementation, dietary patterns, food groups, individual nutrients, fish oil, obesity, dust mites, as well as studies focusing on maternal diet or asthma genotypes, were considered to be irrelevant.

2.6 | Selection of studies

MP and CI initially screened the titles, abstracts, or descriptors of all publications retrieved by the search and duplicates removed.




FIGURE 1 Flow chart of search details

When there was insufficient information in the abstract to warrant exclusion of the article, the full text of the article was retrieved. Full-text papers were then independently appraised by both reviewers for inclusion and details extracted. Discrepancies were resolved by discussion and consensus that led to agreement. The search was supplemented by cross-checking reference and bibliographies of relevant publications, systematic reviews and reviews.

2.7 | Quality assessment and risk of bias

SKS and MP independently rated all publications included in the meta-analysis using a validated quality assessment tool

according to Zaza et al¹⁸ (Data S2). The scientific soundness was rated using 24 validity questions with possible responses of “yes” or “no.” A value of one was assigned to “yes” and zero to “no.” The final score was presented as the percentage of total (maximum) probable scores for each study. Any discrepancies were addressed with mutual agreement between the authors or with a third reviewer (BE). Finally, the scores from both evaluators were averaged and presented as the final score. The observed median study quality score was used as the dividing point between low quality and high quality.¹⁹ We set the benchmark of 70% and above as a high-quality study while below 70% as low quality.

2.8 | Data extraction

Data extracted included authors, year of publication, study name, study design, geographic location, sample size, age range of target population, follow-up, dietary exposure, outcome measures, dietary and respiratory assessment tools, confounders, main findings (risk estimates along with 95% CI and *P*-values).

2.9 | Data synthesis and statistical analysis

Study outcomes were synthesized and categorized into two groups based on the effect on asthma symptoms (improvement or no improvement). Quantitative data were pooled in statistical meta-analysis software using STATA software version 14.1 (Stata Corp, Texas). For the purpose of the meta-analyses, publications assessing "all" types of fish and "fatty fish" as the exposures and regular fish intake, at least once a week were deemed appropriate. Studies were included in the meta-analysis when effect size was expressed as odds ratio (OR) and their 95% confidence intervals (95%CI) were reported or could be calculated. Eligible publications were categorized into two groups based on the common outcome of interest "current asthma" or "current wheeze" as defined by ISAAC.²⁰

For the purpose of the meta-analysis and due to heterogeneity among study designs, subgroup analyses were conducted based on study design, age range, follow-up and type of fish consumed. Cut-offs for age were defined based on children's age range in publications.

Two assessments were conducted for each outcome. One was a combination of all study designs with "all fish or 'fatty fish' exposure" and the other according to study design and similar age group. Heterogeneity was assessed using the standard Chi-square test and I^2 statistic.²¹ Considerable heterogeneity was considered to be an I^2 value >75%²¹ and statistical significance at the 5% level.²² We applied the random-effects model to estimate the pooled ORs and 95% CIs for "current asthma and 'current wheeze'."

3 | RESULTS

3.1 | Electronic search

The literature search identified a total of 1119 potential publications of which 690 remained, after removal of duplicates (Figure 1). No additional citations were found by cross-checking of reference lists. Only one ongoing RCT was identified from international clinical trial registries. Two reviewers (MP, CI) independently screened the 690 eligible papers by scanning titles and abstracts. A total of 656 were excluded based on publication type (136), no abstract (15) or on PICO criteria as they did not examine the effect of fish intake (160) in association with asthma symptoms (308) in children, (37) leaving 34 full-text articles to be assessed for eligibility. Of the remaining 34 potential studies, 11 full texts were deemed to be inappropriate (1 not intervention, 1 not asthma, 6 no full text available and 3 no English translation), leaving a total of 23 relevant studies.

Specifically, 12 cross-sectional,^{10,23-33} 2 case-control^{34,35} and 9 cohort studies^{11,36-43} examining the effect of fish consumption on asthma symptoms in children.

3.2 | Study characteristics

The database search identified 23 studies conducted from 1992 to July 2017 examining the effect of fish consumption on asthma in children. Two were undertaken in Australia,^{33,34} Japan,^{28,32} Spain^{10,25} and Finland^{35,41}; three in Norway^{38,40,42} and the Netherlands^{31,37,43}; four in Sweden^{11,27,36,39}; one in China,³⁰ Italy,²⁶ France,²³ Central Europe²⁹ and the ISAAC study which involved 20 countries globally.²⁴ Collectively, 163 744 children and adolescents up to 15 years old were investigated with sample sizes ranging from 138 to 50 004. A variety of assessment methods were used to assess dietary intake in children. Twenty studies evaluated fish intake using a number of different questionnaires^{10,11,23,24,26,27,29-34,36-43} ranging from one question to questionnaires consisting of limited food groups/items and detailed Food Frequency questionnaires (FFQ) comprising of a wide variety of foods that capture the entire dietary pattern, while two studies used a 3-day food record^{25,35} and one, a diet history questionnaire.²⁸

In all studies, consumption of "all types" of fish (lean and fatty) were investigated, while in six studies the effect of fatty fish intake on asthma symptoms was measured.^{28,34,36-38,42} Asthma was diagnosed using a questionnaire,^{10,23-32,34-39,41,43} parent-report,⁴² doctor-diagnosed^{11,37,40} and in one study by bronchial hyperresponsiveness (BHR).³³ The characteristics of included studies are shown in (Table S1). Overview of the literature search will be presented based on study design and age group.

3.2.1 | Cohort studies: 0-12 years of age

The nine cohort studies identified, investigated the effect of early fish intake on asthma in 33, 673 children from birth until 12 years of age.^{11,36-43} Seven studies revealed that early introduction and regular consumption of "all fish" were associated with a reduction in asthma symptoms up to 12 years of age^{11,36-41} and two showed no effect.^{42,43}

Two studies reported that early introduction of "all fish" (at 6-9 months of age) and regular consumption (≥ 1 /week) of "all fish"³⁷ and "fatty fish"³⁸ reduced asthma incidence (OR: 0.72; 95%CI: 0.55-0.93),³⁸ prevalence of wheeze (OR: 0.64; 95% CI: 0.43-0.94)³⁷ and use of medication (OR: 0.75; 95%CI: 0.58-0.96)³⁸ in children up to 4 years of age.^{37,38} Conversely, Kiefte-de-Jong et al noted that no fish during the first 12 months of life and introduction of "all fish" between 0 and 6 months was associated with an increase in prevalence of wheezing [(OR: 1.57; 95%CI: 1.07-2.31), (OR:1.53; 95%CI: 1.07-2.19)], respectively, at 48 months as compared to a reduction in the prevalence of wheezing with fish introduction at age 6-12 months (OR: 0.64; 95% CI: 0.43-0.94).

Beneficial effects of "all fish" intake were also documented in children during the first 5 years of life. Fish consumption

before 9 months had a protective effect for recurrent wheezing (OR: 0.6; 95% CI 0.4-0.8),³⁹ multiple trigger wheeze (OR: 0.6; 95%CI:0.3-0.99),³⁹ episodic viral wheeze (OR:0.6; 95%CI:0.4-0.99),³⁹ asthma risk (OR: 0.73; 95%CI: 0.55-0.97),⁴¹ (OR: 0.84; 95% CI = 0.57-1.22),⁴⁰ all asthma ($P < .001$)⁴¹ and atopic asthma ($P < .05$)⁴¹ up to 5 years as fish intake increased from two to three times per month ($OR_{adj} = 0.82$ 95% CI: 0.54-1.29) to once a week ($OR_{adj} = 0.66$, 95% CI: 0.43-1.01) and at least once a week ($OR_{adj} = 0.55$; 95% CI 0.34-0.87; p trend = 0.003)¹¹ a dose response was observed.

A Swedish cohort which involved 3285 children with follow-up till 12 years of age, regular "all fish" intake ($\geq 2-3$ times/month) at age of 1 year was associated with reduced risk of prevalence (OR_{adj} : 0.71; 95%CI: 0.57-0.87) and incidence (OR_{adj} : 0.80; 95%CI: 0.65-0.98) of asthma up to age 12 years, after adjusting for confounders of parental history of allergic disease, sex, maternal smoking during pregnancy.³⁶ However, no significant association was observed between fish intake at age 8 years and incidence of asthma at 12 years (p trend = 0.303).³⁶ This finding is in agreement with two other cohorts that documented no relationship among early, late or long-term intake of any kind of fish on asthma in children at ages 2⁴² ($P = .16$) and 8 years.⁴³ However, Willers et al⁴³ did report that early fish intake was inversely associated with BHR in children at age 8 years old.

3.2.2 | Case-control studies: 5-15 years of age

A beneficial effect of fish intake on asthma in children was observed in 2 case-control studies.^{34,35} In a nested-cohort study which was a subgroup of the Finnish DIPP study, 182 children with asthma and 728 controls participated and were observed at 3, 6, 12 months and thereby annually up to 6 years.³⁵ Overall, early introduction of "all fish" and fish products (6-8 months) was associated with a reduction in "all asthma" risk (OR_{adj} : 0.87; 95%CI: 0.77-0.98) in children at 6 years adjusting for confounders. Another case-control study investigating the impact of fish intake on asthma symptoms in 584 children aged 8-11 years also reported a protective effect.³⁴ After adjusting for confounders, children who ate fresh, "fatty fish" at least once a week had a significantly reduced risk of current asthma (OR_{adj} : 0.26; 95%CI: 0.09-0.72) as compared to "any fresh" fish (OR_{adj} : 0.52; 95% CI: 0.24-1.15) and "non-fatty" fish (OR_{adj} : 0.68; 95% CI: 0.3-1.54) consumers. No reduction in asthma risk was observed with canned fish consumption.

3.2.3 | Cross-sectional studies: 0-15 years

The literature search retrieved 12 cross-sectional studies undertaken in 128 577 children aged 0-15 years.^{10,23-33} Five studies observed that regular "all fish" intake (>1 /week) was associated with a reduction in lifetime prevalence of asthma (OR_{adj} : 0.92; 95%CI: 0.78-1.08),²⁴ current wheeze [$(OR_{adj}: 0.85; 95\%CI:0.74-0.97)$]²⁴; (OR: 0.44; 95%CI: 0.21-0.93),³¹ current wheeze with atopy positive (OR_{adj} : 0.51; 95%CI: 0.32-0.81),²⁴ doctor-diagnosed asthma

(OR: 0.54; 95%CI: 0.35-0.84),²⁷ night-time breathlessness (OR: 0.36; 95%CI:0.17-0.78),²⁷ current asthma (OR: 0.51; 95%CI: 0.31-0.84),²⁷ (OR: 0.34; 95%CI: 0.13-0.85),³¹ past-year wheeze (OR adj: 0.75; 95%CI: 0.53-0.93),²³ past-year wheeze and atopy negative (OR: 0.61; 95%CI: 0.43-0.87),²³ atopic asthma with BHR (OR:0.12; 95%CI: 0.02-0.66),³¹ atopic wheeze with BHR (OR: 0.15; 95%CI: 0.03-0.63)³¹ and BHR (OR:0.35; 95%CI: 0.1-0.9).³³ While one study²⁹ reported that irregular fish intake (<1 /month) was associated with an increase in "persistent cough" (OR: 1.18; 95%CI:1.04-1.34), "wheeze ever" (OR: 1.14; 95%CI:1.03-1.25) and "current wheeze" (OR: 1.21; 95%CI: 1.06-1.39). A recent study undertaken by Xu et al,³⁰ including 13, 877 children, 0-14 years old, observed that fish and shrimp intake triggered asthma episodes by 14% ($P < .05$), although estimated effect size was lacking. Adverse effects of "all fish" intake were also reported in a Japanese study of 1673 asthmatic children and 22 109 controls aged 6-15 years.³² After adjustment for confounders (age, gender, parental asthma), a statistically significant higher prevalence of asthma was observed in children consuming fish 1-2 times/week (OR_{adj} : 1.117; 95%CI: 1.005-1.241) and more than or equal to 3-4 times/week (OR_{adj} : 1.319; 95%CI: 0.896-1.943) than among those who consumed no fish (OR_{adj} : 1.039; 95%CI: 0.785-1.376).

3.2.4 | Quality assessment

Twenty-one of 23 studies were rated as "high quality." The average quality score was 82.1% with the highest score 100% and the lowest score 64% (Table 1). Most of the cohort studies included in the meta-analysis were of high quality, 8 of 9 studies scored above 90%. The studies that were rated lower quality were due to their failure to control for potential confounders and address possible biases. Using t test, we checked if there were any differences in the quality score for the selected studies between two reviewers and found no major difference in average score (P -value = .33).

3.3 | Meta-analysis

A total of nineteen studies were used in the meta-analysis.^{10,11,23-29,31,32,34-37,39,40,42,43}

3.3.1 | Current asthma

Assessment of cohort studies alone^{11,40,42} revealed a statistical significant effect²¹ of "all fish" intake on "current asthma" in children 0-4 years ($I^2 = 11.5\%$; $P = .32$; (OR: 0.75; 95% CI: 0.60-0.95). Overall, a 25% reduction in "current asthma" was observed with "all fish" intake in children up to 4 years old (Figure 2).

3.3.2 | Current wheeze

A pooled effect for current wheeze and "All fish" intake was found in cohort studies regarding children up to 4.5 years [$I^2 = 0\%$, $P = .809$; (OR: 0.62; 95%CI: 0.48-0.80)] (Figure 3). A 38% reduction in

TABLE 1 Summary table of Quality Assessment of relevant studies in this systematic review (Zaza)¹⁸

Study	Reviewers	Study design	Population Density	Population Description	Time points	Entire sampling or Probability sampling	Eligibility Criteria	Exposure Measures			Outcome Measures			Statistical Test Suitable	Adjustment For Confounders	Bias Discussed	Bias not covered	Overall Average Score (%)
								Validity	Reliability	Validity	Reliability	Validity	Reliability					
Peat et al ³³	SKS/MP	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	67.5%
Hodge et al ³⁴	SKS/MP	✓	✓	✓	✓	×	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	70.5%
Takemura et al ³²	SKS/MP	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	83.0%
Farchi et al ²⁶	SKS/MP	✓	✓	✓	✓	×	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	73.0%
Nafstad et al ⁴⁰	SKS/MP	✓	×	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	91.0%
Kim et al ²⁷	SKS/MP	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	79.5%
Kull et al ¹¹	SKS/MP	✓	×	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	95.0%
Tabak et al ³¹	SKS/MP	✓	×	✓	✓	×	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	75.0%
Chatzi et al ¹⁰	SKS/MP	✓	×	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	82.0%
Antova et al ²⁹	SKS/MP	✓	✓	✓	✓	×	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	75.0%
Nagel et al ²⁴	SKS/MP	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	83.0%
Oien et al ⁴²	SKS/MP	✓	×	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	95.0%
Rodriguez et al ⁵	SKS/MP	✓	×	✓	×	×	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	73.0%
Goksor et al ³⁹	SKS/MP	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	91.0%
Willers et al ⁴³	SKS/MP	✓	×	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	93.0%
Kunitsugu et al ²⁸	SKS/MP	✓	×	✓	✓	×	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	73.0%
Kiefte de Jong et al ³⁷	SKS/MP	✓	×	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	95.0%
Dotterund et al ³⁸	SKS/MP	✓	✓	✓	✓	✓	×	✓	✓	✓	✓	✓	✓	✓	×	✓	✓	80.5%
Magnusson et al ³⁶	SKS/MP	✓	×	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	91.0%
Nwaru et al ⁴¹	SKS/MP	✓	×	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	97.5%
Lumia et al ³⁵	SKS/MP	✓	×	✓	✓	×	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	78.0%
Saadeh et al ²³	SKS/MP	✓	×	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	82.0%
Xu et al ³⁰	SKS/MP	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	×	✓	✓	✓	64.0%

Key: ✓, criteria satisfied; ×, criteria not satisfied.

“current wheeze” was observed with “All fish” intake in children up to 4.5 years old.

3.3.3 | Fatty fish versus No fatty fish

When analysing the effect of “Fatty fish” intake in children aged 8-14 years^{28,34} for “current asthma” in the combined subanalysis, an overall effect was observed [$I^2 = 0\%$, P -value = .481; (OR: 0.35; 95%CI: 0.18-0.67)] (Figure 4). Fatty fish intake reduced “current asthma” in children aged 8-14 years by 65% as compared to “no fish” intake.

In contrast, no association was observed for “all fish” intake on “current asthma” in children aged (2-15 years) when combining all study designs^{11,24,25,27,28,31,32,34-36,40,42,43} (Figure S1) or in subanalysis based on study designs separately (Figure S2). No pooled age-specific effects were also found for “current wheeze” and “All fish” intake in children (0-13 years) irrespective of study design (Figures S3-S5). The same trend was established for “Fatty fish” intake vs “current asthma” in children (0-14 years) in the combined subanalysis (Figure S6).

4 | DISCUSSION

The aim of this qualitative synthesis and meta-analysis was to clarify the role of fish intake on different asthma outcomes in children. Although the meta-analysis produced mixed results, three important findings were highlighted. Firstly, early introduction of “all fish” intake during infancy (6-9 months of age) and regular consumption (at least once/week) of “all fish” reduced “current asthma” by 25% and “current wheeze” by 38% in children up to 4.5 years old.^{11,37,39,40,42} While “fatty fish” intake seems to confer an overall “prophylactic effect” in older children 8-14 years and reduced “current asthma” by 65%.^{28,34}

Summarizing the results of the database search, in 15/23 studies a protective effect was observed between fish consumption on asthma symptoms in children,^{11,23,24,27,29,31,33-41} 6/23 no effect^{10,25,26,28,42,43} and 2/23 an adverse effect.^{30,32} Two important results were highlighted in these observational studies. Early introduction during the first year of life (between 6 and 9 months) and regular consumption of fish (at least once a week) decreased risk, prevalence and asthma symptoms in children up to 14 years of age. This finding is in accordance with other observational studies that have found early introduction and frequent consumption of fish were associated with decreased asthma symptoms and improvement in pulmonary function in children and adolescents.^{9,44-46}

Our meta-analysis of nineteen¹⁹ publications showed an overall beneficial effect of “all fish” intake on “current asthma” and “current wheeze” in children up to 4.5 years old in five cohort studies alone^{11,37,39,40,42} and of “fatty fish” intake on “current asthma” in 8-14 year olds.^{28,34} The prophylactic effects documented in the cohort studies for young children up to 4.5 years old may be explained from research on biological mechanisms that have shown

that, the development and maturation of the immune system starts early in foetal life continue through infancy and early childhood.^{47,48} Infancy is a period in which the immune cells have an increased vulnerability and susceptibility to environmental exposures such as diet. A diet rich in Omega-3 fatty acids and lower Omega-6: Omega-3 fatty acid ratio may result in an increased incorporation of EPA and DHA into cell membranes at the expense of arachidonic acid. More EPA and DHA in the cell membrane result in decreased production of arachidonic acid-derived pro-inflammatory eicosanoids. EPA and DHA act as substrates for COX and LOX enzymes producing anti-inflammatory eicosanoids, protectins, resolvins and maresins that appears to exert anti-inflammatory and inflammatory resolving actions.⁴⁹ Apart from anti-inflammatory properties, Omega-3 fatty acids are able to modulate immune responses by promoting Th 1 cell generation, thereby reducing airway inflammation, improving pulmonary function and decreasing asthma symptoms.¹³

Additional protective effects on asthma symptoms were observed in two studies examining the effect of “fatty fish” as opposed to “all fish” in children aged 8-11 years old.^{34,38} Hodge et al³⁴ reported a 25% reduction in “current asthma” when fatty fish was consumed as compared to lean or no fish in children. Furthermore, no reduction in asthma risk was observed with the consumption of canned and processed fish. Perhaps, processing and food additives may alter the biological activity of Omega-3 fatty acids in fish oils. Dotterud et al³⁸ reported a 32% reduction in incidence of asthma with fatty fish consumption in infants at 2 years old, after adjusting for confounders. A beneficial effect of fatty fish intake was also noted in the studies undertaken by Kunitsugu and Oien, although results were not statistically significant, most probably explained by heterogeneity between study designs and small sample size.^{28,42} Our meta-analysis confirmed an overall “prophylactic” effect of fatty fish intake in older children.^{28,34} A 65% reduction in “current asthma” was observed in children aged 8-14 years.^{28,34}

The findings of this meta-analysis highlight that the type of fish consumed whether fatty or lean may matter. Fatty fish as compared to lean fish is a rich source of Omega-3 fatty acids which are able to counteract the action of Omega-6 fatty acids metabolites by down-regulating pro-inflammatory and immunological pathways, thereby preventing asthma development.^{50,51} Another possible explanation why fish consumption reduced asthma risk in children as compared to null effects reported from fish oil supplementation during childhood⁵² is that regular fish consumption may be a proxy for other healthy lifestyle habits that might promote the beneficial effect. In addition, fresh fish might have better bioavailability and absorption rate as opposed to fish oil. And apart from EPA/DHA, fresh fish contains other bioactive molecules such as selenium, iodine, vitamin D, potassium and B-vitamins that might interact synergistically providing these prophylactic properties.⁵³

On the other hand, our meta-analysis showed no statistical significant results for 14 studies (10 cross-sectional, 2 cohorts and 2 case-controls) on “current asthma” or “current wheeze” in children 6-15 years old.^{10,23-29,31,32,34,35,42,43} A possible reason for




FIGURE 2 Forest plot of cohort studies children (0-4 years old) for “All fish” intake versus “No fish” and outcome “current asthma” [Colour figure can be viewed at wileyonlinelibrary.com]




FIGURE 3 Forest plot of cohorts comparing the effect of “All fish” intake versus “No fish” on “current wheeze” in children aged 0-4.5 years [Colour figure can be viewed at wileyonlinelibrary.com]




FIGURE 4 Forest plot in the combined analysis comparing “Fatty Fish” intake versus “No Fatty Fish” in children (8-14 years) for “current asthma” [Colour figure can be viewed at wileyonlinelibrary.com]

the null effects observed might be due to heterogeneity among study methodologies which included: population differences, sample size, children’s age, gender, asthma definition, exposure and outcome measures, possible confounders, design of dietary questionnaires and food frequency categories, amount or type of fish consumed (fatty versus lean), quantity EPA/DHA, fish consumption patterns since nutritional guidelines may vary in

each country and food preparation methods. Use of invalidated and self-administered food frequency questionnaires, 3-day diet history for the recording of dietary intake require education and expertise are prone to information bias and response errors.^{43,28} Even though data on habitual diet were collected retrospectively from parents, in young children, it is well-established that parents provide reliable information on children’s diet.⁵⁴ Misclassification

of asthma may have occurred in young children, since the diagnosis of asthma is uncertain in children younger than 5 years and wheezing is often the result of respiratory infection.^{55,56} Furthermore, spirometry which is considered to be the "gold standard" of pulmonary function testing in the diagnosis of asthma, cannot be performed efficiently in children younger than 5 years and diagnosis of asthma in young children is based on parents' report of symptoms.⁵⁵ In addition, parental allergic disease and early onset of allergic disease in children may cause parents to delay introduction or to avoid fish in the child's diet, thus influencing asthma outcome.

Two cross-sectional studies documented increased risk of asthma prevalence in children consuming fish in Japan and China.^{30,32} Today, the eating patterns of Japanese and Chinese children have changed to a more Westernized diet.^{57,58} Processed^{23,59} and pickled foods⁶⁰ as well as a high salt intake^{61,62} have been associated with an increase in bronchial hyperresponsiveness and asthma symptoms in children. In both studies confounding factors such as overweight, socio-economic level, maternal education, type of fish consumed (lean vs fatty), poor adherence to asthma therapy were not included in regression models and may have contributed to the adverse effects.

One of the strengths of this systematic review is the extensive literature search and large number of recent high-quality publications included which increases the power of the analysis. There are very few meta-analysis in the literature that examine fish intake and asthma in children (Yang, 2013; Zhang, 2017)^{14,63} and recent reviews (Best, 2016; Miles, 2017)^{12,13} address Omega-3 fatty acid intake during pregnancy. Our meta-analysis focuses exclusively on children and adolescents and adds to the existing evidence. Another strong point is the inclusion of cohort studies in the meta-analysis which are considered robust and provide strong scientific evidence because they measure events in temporal sequence thereby distinguishing causes from effects.⁶⁴ Another drawback might be publication bias where studies reporting no association between fish intake and asthma may not be published by periodicals.

5 | CONCLUSION

We conducted an up-to-date systematic search to determine the impact of fish intake in childhood asthma. This meta-analysis suggests that introduction of fish early in life (6-9 months) and regular consumption of all fish (at least once a week) reduces asthma and wheeze in children up to 4.5 years old, while fatty fish intake may be beneficial in older children. Future well-designed clinical trials are recommended to confirm the promising findings documented in this literature analysis.

CONFLICT OF INTEREST

The authors declare no potential conflicts of interest.

AUTHOR CONTRIBUTION

MP and CI jointly conducted searches, assessed inclusion, extracted data and assessed validity. MP drafted the manuscript and conducted the meta-analysis. BE and SKS made substantial contributions to analysis and interpretation of data quality. All authors critically reviewed the manuscript and approved the final version as submitted.

ORCID

M. M. Papamichael  <http://orcid.org/0000-0003-1169-7141>

B. Erbas  <http://orcid.org/0000-0001-9597-418X>

REFERENCES

1. Asher M, Montefort S, Björkstén B, et al. Worldwide time trends in the prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and eczema in childhood: ISAAC Phases One and Three repeat multicountry cross-sectional surveys. *Lancet*. 2006;368:733-743.
2. Lai C, Beasley R, Crane J, Foliaki S, Shah J, Weiland S. The ISAAC Phase Three Study Group: global variation in the prevalence and severity of asthma symptoms: Phase Three of the International Study of Asthma and Allergies in Childhood (ISAAC). *Thorax*. 2009;64:476-483.
3. Devereux G. Session 1: Allergic disease Nutrition as a potential determinant of asthma. *Proc Nutr Soc*. 2010;69:1-10.
4. Simopoulos AP. Omega-6/omega-3 essential fatty acid ratio and chronic diseases. *Food Rev Int*. 2004;20:77-90.
5. Simopoulos AP. The importance of the ratio of omega-6/omega-3 essential fatty acids. *Biomed Pharmacother*. 2002;56:365-379.
6. Calder PC. Omega-3 polyunsaturated fatty acids and inflammatory processes: nutrition or pharmacology? *Br J Clin Pharmacol*. 2013;75:645-662.
7. Calder PC. Marine omega-3 fatty acids and inflammatory processes: effects, mechanisms and clinical relevance. *Biochim Biophys Acta*. 2015;1851:469-484.
8. Thien FC, De Luca S, Woods R, Abramson MJ. Dietary marine fatty acids (fish oil) for asthma in adults and children. *Cochrane Database Syst Rev*. 2010;2:CD001283.
9. Alm B, Goksör E, Thengilsdottir H, et al. Early introduction of fish decreases the risk of preschool wheezing disorder. *Allergy Eur J Allergy Clin Immunol*. 2009;64:64-65.
10. Chatzi L, Torrent M, Romieu I, et al. Diet, wheeze, and atopy in school children in Menorca, Spain. *Pediatr Allergy Immunol*. 2007;18:480-485.
11. Kull I, Bergström A, Lilja G, Pershagen G, Wickman M. Fish consumption during the first year of life and development of allergic diseases during childhood. *Allergy*. 2006;61:1009-1015.
12. Best KP, Gold M, Kennedy D, Martin J, Makrides M. Omega-3 long-chain PUFA intake during pregnancy and allergic disease outcomes in the offspring: a systematic review and meta-analysis of observational studies and randomized controlled trials. *Am J Clin Nutr*. 2016;103:128-143.
13. Miles EA, Calder PC. Can early Omega-3 fatty acid exposure reduce risk of childhood allergic disease? *Nutrients*. 2017;9:pii: E784.
14. Yang H, Xun P, He K. Fish and fish oil intake in relation to risk of asthma: a systematic review and meta-analysis. *PLoS ONE*. 2013;8:e80048.
15. Moher D, Liberati A, Tetzlaff J, Altman DG, Group TP. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.

16. Liberati A, Altman D, Tetzlaff J, Mulrow C, Gøtzsche P, Ioannidis J. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med.* 2009;6:e1000100.
17. GINA. 2017 Pocket Guide for Asthma Management and Prevention: for adults and children older than 5 years. 2017.
18. Zaza S, Wright-De Agüero LK, Briss PA, et al. Data collection instrument and procedure for systematic reviews in the Guide to Community Preventive Services. *Am J Prev Med.* 2000;18:44-74.
19. Brouwers MC, Johnston ME, Charette ML, Hanna SE, Jadad AR, Browman GP. Evaluating the role of quality assessment of primary studies in systematic reviews of cancer practice guidelines. *BMC Med Res Methodol.* 2005;5:8.
20. Asher M, Keil U, Anderson H, et al. International Study of Asthma and Allergies in Childhood (ISAAC): rationale and methods. *Eur Respir J.* 1995;8:483-491.
21. Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011].* The Cochrane Collaboration, 2011. Available from <http://handbook.cochrane.org>.
22. Egger M, Smith GD, Phillips AN. Meta-analysis: principles and procedures. *Br Med J.* 1997;315:1533.
23. Saadeh D, Salameh P, Caillaud D, et al. Prevalence and association of asthma and allergic sensitization with dietary factors in schoolchildren: data from the french six cities study. *BMC Public Health.* 2015;15:993.
24. Nagel G, Weinmayr G, Kleiner A, Garcia-Marcos L, Strachan D, Group tiPTS. Effect of diet on asthma and allergic sensitisation in the International Study on Allergies and Asthma in Childhood (ISAAC) Phase Two. *Thorax.* 2010;65:516-522.
25. Rodríguez-Rodríguez E, Perea J, Jiménez A, Rodríguez-Rodríguez P, López-Sobaler A, Ortega R. Fat intake and asthma in Spanish schoolchildren. *Eur J Clin Nutr.* 2010;64:1065-1071.
26. Farchi S, Forastiere F, Agabiti N, et al. Dietary factors associated with wheezing and allergic rhinitis in children. *Eur Respir J.* 2003;22:772-780.
27. Kim JL, Elfman L, Mi Y, Johansson M, Smedje G, Norback D. Current asthma and respiratory symptoms among pupils in relation to dietary factors and allergens in the school environment. *Indoor Air.* 2005;15:170-182.
28. Kunitsugu I, Okuda M, Murakami N, et al. Self-reported seafood intake and atopy in Japanese school-aged children. *Pediatr Int.* 2012;54:233-237.
29. Antova T, Pattenden S, Nikiforov B, et al. Nutrition and respiratory health in children in six Central and Eastern European countries. *Thorax.* 2003;58:231-236.
30. Xu D, Wang Y, Chen Z, et al. Prevalence and risk factors for asthma among children aged 0-14 years in Hangzhou: a cross-sectional survey. *Respir Res.* 2016;17:1-8.
31. Tabak C, Wijga AH, de Meer G, Janssen NA, Brunekreef B, Smit HA. Diet and asthma in Dutch school children (ISAAC-2). *Thorax.* 2006;61:1048-1053.
32. Takemura Y, Sakurai Y, Honjo S, et al. The relationship between fish intake and the prevalence of asthma: the Tokorozawa childhood asthma and pollinosis study. *Prev Med.* 2002;34:221-225.
33. Peat J, Salome C, Woolcock A. Factors associated with bronchial hyperresponsiveness in Australian adults and children. *Eur Respir J.* 1992;5:921-929.
34. Hodge L, Salome C, Peat J, Haby M, Xuan W, Woolcock A. Consumption of oily fish and childhood asthma risk. *Med J Aust.* 1996;164:137-140.
35. Lumia M, Takkinen HM, Luukkainen P, et al. Food consumption and risk of childhood asthma. *Pediatr Allergy Immunol.* 2015;26:789-796.
36. Magnusson J, Kull I, Rosenlund H, et al. Fish consumption in infancy and development of allergic disease up to age 12 y. *Am J Clin Nutr.* 2013;97:1324-1330.
37. Kieft-de Jong JC, de Vries JH, Franco OH, et al. Fish consumption in infancy and asthma-like symptoms at preschool age. *Pediatrics.* 2012;130:1060-1068.
38. Dotterud CK, Storrø O, Simpson MR, Johnsen R, Øien T. The impact of pre-and postnatal exposures on allergy related diseases in childhood: a controlled multicentre intervention study in primary health care. *BMC Public Health.* 2013;13:1.
39. Goksör E, Alm B, Thengilsdottir H, Pettersson R, Åberg N, Wennergren G. Preschool wheeze-impact of early fish introduction and neonatal antibiotics. *Acta Paediatr.* 2011;100:1561-1566.
40. Nafstad P, Nystad W, Magnus P, Jaakkola JJ. Asthma and allergic rhinitis at 4 years of age in relation to fish consumption in infancy. *J Asthma.* 2003;40:343-348.
41. Nwaru BI, Takkinen H-M, Niemelä O, et al. Timing of infant feeding in relation to childhood asthma and allergic diseases. *J Allergy Clin Immunol.* 2013;131:78-86.
42. Oien T, Storro O, Johnsen R. Do early intake of fish and fish oil protect against eczema and doctor-diagnosed asthma at 2 years of age? A cohort study. *J Epidemiol Community Health.* 2010;64:124-129.
43. Willers SM, Wijga AH, Brunekreef B, et al. Childhood diet and asthma and atopy at 8 years of age: the PIAMA birth cohort study. *Eur Respir J.* 2011;37:1060-1067.
44. D'Innocenzo S, Matos S, Prado MS, et al. Dietary pattern, asthma, and atopic and non-atopic wheezing in children and adolescents: SCAALA study, Salvador, Bahia State, Brazil. *Cadernos de Saúde Pública.* 2014;30:1849-1860.
45. Ellwood P, Asher M, Björkstén B, Burr M, Pearce N, Robertson C. Diet and asthma, allergic rhinoconjunctivitis and atopic eczema symptom prevalence: an ecological analysis of the International Study of Asthma and Allergies in Childhood (ISAAC) data. *Eur Respir J.* 2001;17:436-443.
46. Arvaniti F, Priftis KN, Papadimitriou A, et al. Adherence to the Mediterranean type of diet is associated with lower prevalence of asthma symptoms, among 10-12 years old children: the PANACEA study. *Pediatr Allergy Immunol.* 2011;22:283-289.
47. Noutsios GT, Floros J. Childhood asthma: causes, risks, and protective factors; a role of innate immunity. *Swiss Med Wkly.* 2014;144:w14036.
48. Prescott SL. Early-life environmental determinants of allergic diseases and the wider pandemic of inflammatory noncommunicable diseases. *J Allergy Clin Immunol.* 2013;131:23-30.
49. Galli C, Calder PC. Effects of fat and fatty acid intake on inflammatory and immune responses: a critical review. *Ann Nutr Metab.* 2009;55:123-139.
50. Kremmyda L-S, Vlachava M, Noakes PS, Diaper ND, Miles EA, Calder PC. Atopy risk in infants and children in relation to early exposure to fish, oily fish, or long-chain omega-3 fatty acids: a systematic review. *Clin Rev Allergy Immunol.* 2011;41:36-66.
51. Moreno-Macias H, Romieu I. Effects of antioxidant supplements and nutrients on patients with asthma and allergies. *J Allergy Clin Immunol.* 2014;133:1237-1244.
52. Muley P, Shah M, Muley A. Omega-3 fatty acids supplementation in children to prevent asthma: is it worthy?-A systematic review and meta-analysis. *J Allergy.* 2015;2015:312052.
53. Weichselbaum E, Coe S, Buttriss J, Stanner S. Fish in the diet: a review. *Nutr Bull.* 2013;38:128-177.
54. Livingstone M, Robson P. Measurement of dietary intake in children. *Proc Nutr Soc.* 2000;59:279-293.
55. Katsardis C, Koumbourlis A, Anthracopoulos M, Paraskakis E. *Paediatric Pulmonary Function Testing.* New York, USA: Nova; 2015.
56. Lasso-Pirot A, Delgado-Villalta S, Spanier AJ. Early childhood wheezers: identifying asthma in later life. *J Asthma Allergy.* 2015;8:63-73.
57. Murata M. Secular trends in growth and changes in eating patterns of Japanese children. *Am J Clin Nutr.* 2000;72:1379s-1383s.

58. Zhang J, Wang H, Wang Y, et al. Dietary patterns and their associations with childhood obesity in China. *Br J Nutr*. 2015;113:1978-1984.
59. Patel S, Custovic A, Smith JA, Simpson A, Kerry G, Murray CS. Cross-sectional association of dietary patterns with asthma and atopic sensitization in childhood - in a cohort study. *Pediatr Allergy Immunol*. 2014;25:565-571.
60. Akcay A, Tamay Z, Hocaoglu AB, Ergin A, Guler N. Risk factors affecting asthma prevalence in adolescents living in Istanbul, Turkey. *Allergol Immunopathol*. 2014;42:449-458.
61. Arvaniti F, Priftis K, Papadimitriou A, et al. Salty-snack eating, television or video-game viewing, and asthma symptoms among 10- to 12-year-old children: the PANACEA study. *J Am Diet Assoc*. 2011;111:251-257.
62. Corbo G, Forastiere F, De Sario M, et al. Wheeze and asthma in children: associations with body mass index, sports, television viewing, and diet. *Epidemiology*. 2008;19:747-755.
63. Zhang GQ, Liu B, Li J, et al. Fish intake during pregnancy or infancy and allergic outcomes in children: a systematic review and meta-analysis. *Pediatr Allergy Immunol*. 2017;28:152-161.
64. Song JW, Chung KC. Observational studies: cohort and case-control studies. *Plast Reconstr Surg*. 2010;126:2234-2242.

Websites:

Australian and New Zealand Clinical trial registry: <http://www.anzctr.org.au/trialSearch.aspx>
International Clinical trials registry: <http://www.clinicaltrials.gov>
European Union clinical trials registry: <http://www.clinicaltrialsregister.eu>

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

How to cite this article: Papamichael MM, Shrestha SK, Itsiopoulos C, Erbas B. The role of fish intake on asthma in children: A meta-analysis of observational studies. *Pediatr Allergy Immunol*. 2018;29:350-360. <https://doi.org/10.1111/pai.12889>